

T. (*Erythrosperma*) hispanicum n. sp. — Humile, 4—10 cm altum, gracile; folia læte viridia supra subglabra vel pilis tenuibus sparsis prædita, subtus tantum in nervo sparse araneosa, petiolis angustis, non coloratis vel rarius ± vinoso-purpureis, lamina in foliis exterioribus lobis retroversis, arcuatis, subobtusis, subintegris, in foliis interioribus lobis triangularibus acutis, subpatulis, integris vel parce dentatis; scapi 1—plures, parce araneosi, in statu exsiccato 2—2.5 mm crassi; involucrium pruinosum, 10—14 mm longum, c. 8 mm latum; squamæ exteriores patulæ, tenues, angustissime albo-marginatæ, ± violascentes, ecorniculatæ, 1.5—2 mm latæ, c. 6 mm longæ, squamæ interiores apice suberosæ, atroviolaceæ, ecorniculatæ; ligulæ luteæ, marginales stria atroviridi ornata; antheræ polliniferæ; stigmata atroviridia; achenium rubiginosum, 2.5 mm longum, 0.8 mm latum, superne acute spinulosum, ceterum obtuse tuberculatum vel sublæve, sensim in pyramidem 0.7 mm longam, cylindricam abiens, rostro 7—8 mm longo, pappo albo. — H., Aranjuez prope urbem Madrid, in graminosis juxta stationem viæ ferratæ.

Icon.: Fig. 44 f, 47 e.

T. (*Erythrosperma*) marginellum n. sp. — Humile, c. 12 cm altum, gracile; folia obscure viridia, subglabra, petiolis longis angustis, inferne ± vinoso-purpureis, lamina lobis brevibus, patulis, subobtusis, integris, vel in margine superiore dentibus parvis paucis præditis, interlobiis manifeste dentatis; scapi plures, leviter areneosi, in statu exsiccato c. 3 mm crassi; involucrium 10—12 mm longum, pruinosum; squamæ exteriores breves, ecorniculatæ, late albomarginatæ, c. 4 mm longæ, 2—3 mm latæ, marginibus breviter araneosis, subpatulæ — subadpressæ, squamæ interiores cornibus parvis præditæ; ligulæ luteæ, marginales subtus stria lata, atroviridi ornata; antheræ polliniferæ; stigmata obscure viridia; achenium obscure rubiginosum, 3 mm longum, 1 mm latum, superne acute spinulosum, ceterum vulgo ± tuberculatum, raro sublæve, sensim in pyramidem 0.8 mm longam, cylindricam abiens, rostro 4 mm longo, pappo albo. — H., Aranjuez prope urbem Madrid, in graminosis juxta stationem viæ ferratæ.

Icon.: Fig. 44 b, 47 s.

T. maroccanum n. sp. — Humile, 6—15 cm altum, gracile; folia læte viridia, subglabra, tenuia, petiolis angustis, non coloratis vel rarius paullo vinoso-purpureis, lamina lobis triangularibus, acutis, patulis vel paullo retroversis, in margine superiore dentibus paucis, vulgo sat magnis præditis, lobo terminali vulgo triangulari, lobulo apicali et lobulis lateralibus acutis; scapi plures, glabri, 2—4 mm crassi; involucrium 12—15 mm longum, fere epruinosum; squamæ exteriores angustæ, tenues, subpatulæ, anguste albo-marginatæ, 6—7 mm longæ, 1.5—2 mm latæ, marginibus parce araneosis, ecorniculatæ vel raro cornibus parvis præditæ, squamæ interiores apice nigro, angusto, ecorniculato vel raro cornu parvo; ligulæ luteæ, marginales subtus stria atroviridi ornata; antheræ polliniferæ; stigmata viridi-lutea; achenium pallide cinnamomeum, 2.5—2.8 mm longum, 0.8 mm latum, superne acute breviterque spinulosum, ceterum læve, sat sensim in pyramidem 0.5—0.8 longam, subconicam vel subcylindricam abiens, rostro 6—8 mm longo, pappo albo. — M., Atlas major, Amismiz, in declivi in valle supra pagum, c. 1.500 m. In convalle fluminis Ait Messane, in declivi graminoso, humido juxta flumen supra pagum Arround, c. 2.100 m.

Icon.: Fig. 46 e, 47 s Arround; 46 f, 47 7, Amismiz.

T. (*Vulgaria*) messanense n. sp. — Sat robustum, c. 20 cm altum; folia læte viridia, subglabra, tenuia, lata, 3—4 cm lata, petiolis sat brevibus, angustis, nervo mediano supra ± vinoso-purpureo, lamina lobis latis triangularibus, patulis, integris, non profundis; scapi plures, leviter araneosi, in statu exsiccato 3 mm crassi; involucrium 15 mm altum, subpruinosum; squamæ exteriores tenues, patulæ, 2 mm latæ, c. 9 mm longæ, obsolete marginatæ, ecorniculatæ; ligulæ luteæ, dentibus subtus ± roseis, marginales subtus sat obsolete obscure striatæ; antheræ pol-